



Nutritional aspects of non-prey foods in the life histories of predaceous Coccinellidae

Jonathan G. Lundgren*

USDA-ARS, North Central Agricultural Research Laboratory, Brookings, SD 57006, USA

ARTICLE INFO

Article history:

Received 18 December 2008

Accepted 25 May 2009

Available online 23 June 2009

Keywords:

Biological control

Fungus

Lady beetles

Mixed diets

Meta-analysis

Omnivory

Pollen

Polyphagy

Sugar

ABSTRACT

Non-prey foods are an integral component of the diets of most predaceous coccinellids. Under field conditions, numerous coccinellids consume nectar, honeydew, pollen, fruit, vegetation, and fungus. These non-prey foods are used by coccinellids to increase survival when prey is scarce, reduce mortality during diapause, fuel migration, and enhance reproductive capacity. Each of these non-prey foods has unique nutritional and defensive characteristics that influence its suitability for lady beetles. Quantitatively, nutrient and energy contents of these foods are often competitive with, or even exceed that, present in prey. Meta-analyses of literature were used to assess whether (1) some non-prey foods and prey are equivalent foods for coccinellids, and (2) prey-only diets and mixed diets involving prey and non-prey foods are equally suitable for coccinellids. Response variables were categorized as larval performance (e.g., development time, weight at eclosion), adult performance (e.g., adult longevity, weight change), and reproduction (e.g., fecundity, oviposition period). The analyses revealed that pollen is inferior to prey for supporting larval and adult performance, but that adult performance in sugar-fed adults was equal to that of prey-fed adults (although sugar alone does not support reproduction). Larval performance was enhanced substantially when they were reared on mixed diets compared to prey-only diets. Adding sugar to mixed diets strongly improved adult performance and reproduction over prey-only diets, but this was not the case with pollen in mixed diets. These results suggest that coccinellid larvae have more stringent nutritional requirements than adults, and that non-prey foods provide unique nutrients that enhance prey-only diets. Moreover, it suggests that simple carbohydrates are important dietary constituents capable of enhancing both adult performance and reproduction, although it is important to note that a range of prey species of variable quality for coccinellids are evaluated in this database. The literature review presented here suggests that non-prey foods are a critical component of coccinellid nutritional ecology, and may influence the success of conservation biological control programs.

Published by Elsevier Inc.

1. Introduction

The traditional approach to defining the trophic placement of an organism is narrowly defined, and often is based on an organism's primary function within a food web (e.g., herbivore, detritivore, and predator). Lady beetles are no exception to this trend, and are often defined as predators (but see Giorgi et al., 2009; Sutherland and Parrella, 2009). Within the predatory clades, lady beetles are sometimes further partitioned as specialists on aphids, mites, scales, etc. These designations have utility, both in terms of their application to biological control of agronomic pests and when one considers the importance of these prey groups to the life his-

tories of individual lady beetle species. Nevertheless, it is important that the full dietary breadth of an organism be recognized, especially for those species involved in the biological control of pests.

In 1962, Ivo Hodek defined foods of predators as essential and alternative; essential foods support reproduction and development, and alternative foods do not (Hodek and Honěk, 1996). Since this basic distinction was made, it has often been presumed by scientists and biological control practitioners that coccinellids spend their time seeking out their respective essential foods. But nowhere in his initial definition of these terms does Hodek define essential foods as *optimal* foods for maximizing rates of growth, development and reproduction of a predator. This is underscored by the large number of studies that show that different essential foods vary in their suitability for supporting coccinellid fitness (even intraspecifically), and the fact that mixing different prey usually leads to higher fitness in the predator than eating a single prey

* Tel.: +1 605 693 5211; fax: +1 605 693 5240.

E-mail address: Jonathan.Lundgren@ars.usda.gov

item alone (Evans, 2009; Hodek and Honěk, 2009). Moreover, combining several alternative foods can ultimately lead to diets that support reproduction and development of a coccinellid species (Evans, 2000).

Non-prey foods are an important component of most coccinellid diets, and although they receive less attention than prey in the scientific literature, these foods fundamentally shape the life histories of many coccinellid species. Nectar, honeydew, pollen, fruit, vegetation, and fungus are all regularly consumed by coccinellids under field conditions. Given that alternative foods such as these seldom support critical life functions, it is easy to dismiss them as inferior to essential foods in nutrition. An objective review of the literature shows that each of these non-prey foods represents a rich source of nutrients and energy, and a meta-analysis of published studies reveals that non-prey foods are often an integral component of optimal diets for coccinellids.

2. Characteristics of non-prey foods and their function in the life histories of coccinellids

Although it has long been recognized that natural enemies rely on non-prey foods as part of their diet (Forbes, 1881, 1883), synthesis on the importance of these nutritional resources for natural enemies (including coccinellids) has only come about within the last 22 years. Hagen (1986) was one of the first to comprehensively review the role of plant-based foods in the nutrition of natural enemies. Wäckers (2005) defined several traits that affect the suitability of plant-based foods for natural enemies, including their availability, apparency, accessibility, nutritional composition/suitability, and foraging risks associated with each non-prey food. Most recently, Lundgren (2009) provided a comprehensive discussion on the physiological, morphological, nutritional and environmental factors that mediate the interactions among natural enemies and non-prey foods. Within the context of this review, non-prey foods aid in the survival of coccinellids (particularly when prey is scarce or of poor quality), but also support diapause, dispersal, and reproduction of these important natural enemies.

2.1. Nectar

Floral and extrafloral nectars differ from each other in their nutrition and defensive properties. These differences are driven in large part by the distinct functions that floral and extrafloral nectars play in the life histories of the plants that produce them (Bentley, 1977). Floral nectar plays a crucial role in attracting (often very specific) pollinators and facilitates outbreeding in plants. As such, it is guarded against floral larceny by non-pollinators (like coccinellids). The function of extrafloral nectar (EFN) is to attract entomophagous natural enemies of herbivores (like coccinellids), essentially acting as an extension of the plant's defense response to herbivory (Heil et al., 2001; Heil, 2004; Choh et al., 2006; Kost and Heil, 2006). EFN is produced when the plant needs defense from herbivores the most, particularly during periods of vegetative growth. Under most circumstances, EFN is available for a much greater duration than flower-bound nectar. For these reasons, a review of the literature reveals that EFN is fed upon by coccinellids more frequently than floral nectar.

2.1.1. Nectar nutrition

At first glance, nectar is simply a source of sugar— it is primarily composed of sucrose, glucose, and fructose (Percival, 1961; Elias and Gelband, 1975; Petanidou et al., 1996; Heil et al., 2000; Wäckers, 2001; Lundgren, 2009). A closer examina-

tion reveals that a number of mono- and di-saccharides (as well as some oligo-saccharides) are present in many nectars, but at much lower concentrations than sucrose and its metabolites (Torres and Galletto, 2002; Petanidou, 2005). At least 18 sugars have been found in floral nectars, and 15 from EFN (Lundgren, 2009).

Other nutrients, when they occur, are found in much lower quantities. Amino acids give floral nectar its taste and are important in encouraging pollinator fidelity (Baker and Baker, 1977; Gardener and Gillman, 2002; Petanidou et al., 2006). EFN also frequently possesses amino acids (Hanney and Elmore, 1974; Baker et al., 1978; Smith et al., 1990), perhaps to attract specific beneficials. Lipids are reported (infrequently) in nectars (Keeler, 1977; Baker and Baker, 1983; Caldwell and Gerhardt, 1986), as are vitamins (Baker and Baker, 1983; Jakubska et al., 2005). The amount and types of nutrients found in nectar of a particular species depends on many factors, including plant genetics, floral architecture, physiological status of the plant, age of the bloom and previous feeding from it, and environmental conditions including soil (Shuel, 1955; Cruden et al., 1983; Gottsberger et al., 1984). Usually, sugar concentrations from floral and extrafloral nectars are approximately 20% w/v (Baker, 1975; Guerrant and Fiedler, 1981; Adler, 2000).

2.1.2. Nectar defenses

The defenses that discourage nectar thievery from flowers can be categorized as architectural or chemical in nature. The position of the nectary within the flower has substantial influence on the number of insect species that feed on it. Except for the smallest of species, coccinellids (like other large entomophages; Tooker and Hanks, 2000; Galletto and Bernardello, 2004; Vattala et al., 2006) are largely excluded from floral nectar when it occurs deep within the corolla. Abundant stigmas or styles also inhibit some insects from accessing floral nectars. A wide range of secondary chemicals are found in floral nectar (Baker and Baker, 1978; Adler and Irwin, 2005). These secondary chemicals (which include phenolics, alkaloids, glycosides, saponins, non-protein amino acids, alcohols, and ammonia) deter or intoxicate many insects, play an important part in keeping pollinators that have adapted to feed on them loyal, and generally increase the likelihood of successful pollination of the plant (Adler, 2000; Kessler et al., 2008).

Because its role is to attract whatever beneficial arthropods reside in a given habitat, EFN is largely unprotected from nectar-feeding insects. Even so, a few secondary chemicals have been isolated from EFN (Keeler, 1977; Baker and Baker, 1978), but their role is not well understood.

2.1.3. Coccinellids that feed on nectar

Most entomophagous arthropods will feed on sugar if given the chance, and coccinellids are no exceptions. Few observations of flower-visiting coccinellids have irrefutably substantiated nectarivory (Bugg, 1987; Nalepa et al., 1992; Spellman et al., 2006). In contrast, coccinellids are often some of the most frequent visitors to extrafloral nectaries (EFNs) under field conditions (Putman, 1955, 1963; Banks, 1957; Keeler, 1978; Stephenson, 1982; Ricci et al., 2005). This is particularly so during the spring in temperate regions, when EFNs are most abundant on developing vegetation (Ewing, 1913; Watson and Thompson, 1933; Rockwood, 1952), and coccinellids frequently find themselves prey-limited. Pemberton and Vandenberg (1993) present a list of 41 coccinellid species found feeding at the EFNs of dozens of plant species from 15 families. Little has been added to this extensive review of EFN-coccinellid interactions during the past 16 years, except to further substantiate the place of coccinellids as some of the more frequent visitors to EFNs.

The sugars found in nectars are an easily digestible and high energy food, and can dramatically increase survival of coccinellids in the absence of prey (Geyer, 1947; Ibrahim, 1955; Matsuka et al., 1982; Dreyer et al., 1997). It is worth mentioning that not all mono- and oligo-saccharides are equally suitable for coccinellids; clear preferences were displayed by *Harmonia axyridis* Pallas larvae for simple carbohydrates ($n = 13$ sugars tested) under choice conditions (Nijima et al., 1997). Sugar-feeding also supports the flight capacity of coccinellids (Nedved et al., 2001). Finally, although sugar-feeding seldom supports reproduction in coccinellids on its own, sugar consumption can shorten pre-oviposition periods of coccinellids (Smith and Krischik, 1999), and help females to survive reproductive diapause (Hagen, 1962; Reznik and Vaghina, 2006). In the latter case, reductions in prey density initiate a physiological shift in some coccinellids; essentially a trophically induced reproductive diapause. Nutritional resources are shifted from reproduction to fat storage, and sugar resources can help to increase survival and reduce the resorption of eggs during these periods of reproductive diapause (Reznik and Vaghina, 2006).

2.2. Honeydew

Honeydew is a sugary exudate originating from phloem- and xylem-feeding hemipterans (and some lepidopterans), and it differs from nectars in subtle but important ways. Although honeydew production likely began simply as a sugary excretory product that necessarily accompanies the consumption of nutritionally simple plant sap, it also is a clear indicator to predators of the prey insects that produce it. The valuable sugar secreted by sternorrhynchans has since fueled mutualistic associations with ants, which deter predators such as coccinellids from attacking herbivores (Majerus et al., 2007). These complex interactions notwithstanding, honeydew is ubiquitous in most terrestrial habitats (Way, 1963; Downes and Dahlem, 1987), and is a valuable food source that is consumed by numerous coccinellid species.

2.2.1. Honeydew nutrition

Like the original ingested phloem contents, excreted honeydew is primarily comprised of sugars, especially glucose, fructose, and sucrose (Ewart and Metcalf, 1956; Mittler, 1958; Lamb, 1959). Unlike many nectars, honeydew also has a large complement of trisaccharides (some of which are found in few other places in nature), that are produced within the herbivore (Ashford et al., 2000). These trisaccharides have several roles that will be discussed below (Section 2.2.2). In addition to sugars, honeydews possess a wide range of amino acids that often match the original phloem contents in quality if not in quantity; usually less than 2% of the dry weight of honeydew is amino acids (Ewart and Metcalf, 1956; DeVries and Baker, 1989; Yao and Akimoto, 2002; Woodring et al., 2006). Some sterols and vitamins are also found in honeydews at low concentrations (Gray, 1952; Lamb, 1959).

2.2.2. Honeydew defenses

Honeydew contains a number of chemicals that may help to protect the herbivore from foraging coccinellids. First, the trisaccharides created by the insect to offset the osmotic differential between the hemipteran haemocoel and the copious amount of fluid ingested (Costa et al., 1999; Woodring et al., 2006) may have a dual function of making honeydew less palatable to natural enemies. In part, this may be nutritional; some sugars are indigestible or undetectable by natural enemies (Nijima et al., 1997; Wäckers, 2000, 2001; Williams et al., 2005). But trisaccharides also make honeydew evaporate more quickly, thereby

reducing the number of natural enemies that can partake of the dry, sugary residue. Other defensive properties of honeydew can come from the host plant itself. Some of the defensive secondary chemicals found in plants are found in the honeydew of sternorrhynchans, and can in some instances make the area surrounding hemipteran colonies (where honeydew accumulates) repulsive to natural enemies (Molyneux et al., 1990; Bristow, 1991).

2.2.3. Coccinellids that feed on honeydew

In spite of the ants and aversive chemicals that guard it, honeydew is frequently encountered and of high energy value; thus it is not surprising that many coccinellids consume it as part of their diet (Simanton, 1916; Tedders and Schaefer, 1994). This sugar source can extend the lives of beetles in the absence of prey (Putman, 1955; Yinon, 1969). Also, when only poor quality prey are available, consumption of honeydew along with it can support modest reproduction in some coccinellids; e.g., *Coccinella transversalis* Fabricius (Evans, 2000).

In addition to being a source of nutrition, honeydew also arrests and intensifies the foraging of coccinellid adults and larvae (Banks, 1957; Carter and Dixon, 1984; van den Meiracker et al., 1990; Han and Chen, 2002; Seagraves, 2009). The presence of honeydew is a clear indication of a local population of prey, and coccinellids that encounter patches of honeydew turn more often, and forage more intensively in the hopes of encountering the honeydew-producing prey. This shift in foraging results in aggregations of coccinellids within hemipteran-infested areas of plants, and can lead to suppression of pests.

2.3. Pollen

One of the most nutritious of non-prey foods consumed by coccinellids is pollen. This is one of the reasons that this non-prey food has been studied more extensively than any other as a dietary component for coccinellid beetles. Given the restrictions of flower-bound resources (described above in Section 2.1.2), anemophilous pollen species are likely a more widely dispersed and apparent food source for lady beetles than zoophilous pollens. During peak anthesis within a given habitat, pollen grains are a significant food source for numerous insects (Cottrell and Yeagan, 1998; Hoheisel and Fleischer, 2007; Lundgren, 2009). Coccinellids are just one of the many organisms that recognize this abundant resource as food, and respond behaviorally to its availability within a habitat.

2.3.1. Pollen nutrition

Pollen is an excellent source of essential nutrients for insects (Stanley and Linskins, 1974). Fructose, glucose, and sucrose (among other simple sugars) are present within most pollens (Roulston and Buchmann, 2000). Starches are the most important storage carbohydrates, generally comprising less than 2.5% of pollen dry weight (Baker and Baker, 1979, 1982; Roulston and Buchmann, 2000). Protein is one of the most abundant nutrients in pollen, typically comprising 12–61% of dry weight (Roulston et al., 2000). All essential amino acids for insect growth are present, and of these proline is typically one of the most abundant (Erhardt and Baker, 1990; Lundgren and Wiedemann, 2004; Carter et al., 2006). Lipids are also common to most pollens, comprising up to 19% of dry weight (Standifer, 1967; Barbier, 1970). While sterols are present in most pollens (Standifer et al., 1968), these precursors to insect hormones are a limiting nutrient for the development of pollen-fed lady beetles (Lundgren, unpublished data). A wide range of vitamins and inorganic minerals are also prevalent in many pollens (Stanley and Linskins, 1974). Pollen is practically devoid of water, and

although this raises its per-unit nutrient value, coccinellids reared solely on pollen invariably require a supplemental source of water (De Clercq et al., 2005; Michaud and Grant, 2005). Finally, pollens vary in their nutrition both intraspecifically and among species (Lundgren and Wiedenmann, 2004; Lundgren, 2009), and as such the nutritional value of pollen for a coccinellid species can change substantially among testing systems.

2.3.2. Pollen defenses

In addition to the floral defenses discussed in Section 2.1.2., pollen has a range of physiological defenses that protect it from pollinivory. Structurally, the rigid pollen exine must be dismantled, either mechanically or chemically, before the internal nutrients of the grain can be accessed (Roulston and Cane, 2000). Also, numerous textures and appendages (hooks, spines, barbs, etc.) to the exine can inhibit pollinivory by lady beetles. As a case in point, *Coleomegilla maculata* DeGeer adults and larvae were killed when enclosed with the hooked grains of *Helianthus annuus* L. under laboratory conditions (Michaud and Grant, 2005).

Pollen can be chemically defended either through deficiency in key nutrients for insects, and/or with defensive secondary compounds. Anti-nutritive qualities of pollen are speculated based on the observation that anemophilous pollen (which does not benefit from insect dispersal) has a different nutritional composition than that of animal-pollinated plants (Baker and Baker, 1979). A comprehensive review of the literature reveals that many anemophilous plants have starchier pollens than zoophilous pollens (and thus have lower energy content), but it remains unclear whether this is a case of convergent evolution (Roulston and Buchmann, 2000; Roulston et al., 2000). Many pollens are conspicuously toxic to insects (Lundgren, 2009). Although toxic agents within pollens are seldom identified, alkaloids and galitoxins are reported in some pollens (Stanley and Linskens, 1974).

2.3.3. Coccinellids that consume pollen

In sum, at least 39 species of entomophagous coccinellids have been recorded as consuming more than 88 species of pollen under laboratory and field conditions (Lundgren, 2009). Pollen is believed to be most important as food when prey is scarce or of low quality, as is evident in the significant number of instances of spring and fall pollinivory, when prey is scarce in most temperate habitats (Conrad, 1959; Solbreck, 1974; Benton and Crump, 1981; Anderson, 1982; Hemptinne and Desprets, 1986; Triltsch, 1997, 1999). Consistent with this pattern, regions that experience a mid-summer dearth of prey experiences a concomitant increase in mid-summer pollinivory by coccinellids (Ewing, 1913; Hagen, 1962; Ricci et al., 2005; Michaud and Qureshi, 2006).

It should not be assumed that just because pollen is sought most intensively when prey is scarce that this non-prey food does not fulfill a vital role in the diet of coccinellids. At least several species can complete development on a diet of pollen alone, although the resulting adults may have lower fitness than those fed prey-based diets (Hukusima and Itoh, 1976; Smith, 1961; Berkvens et al., 2008). Others use pollen resources to fuel migrations (Solbreck, 1974) and survive dormancy (Anderson, 1981; Triltsch, 1999). Moreover, pollen is important in spermatogenesis (Hemptinne and Naisse, 1987) and, particularly when mixed with prey, may help to promote reproduction in some species (Hemptinne and Desprets, 1986; Michaud, 2000; De Clercq et al., 2005; Omkar, 2006; Berkvens et al., 2008).

Without question, pollinivory has been best studied for the highly omnivorous New World coccinellid, *Coleomegilla macula-*

ta. The first description of pollen-feeding in this species was by Stephen Forbes in 1881, who substantiated pollinivory under field conditions using microscopic gut content analysis. This coccinellid can complete its development on a range of pollen species (Smith, 1961; Smith, 1965; Hodek et al., 1978; Hazzard and Ferro, 1991; Riddick and Barbosa, 1998; Michaud and Grant, 2005; Michaud and Jyoti, 2008;). Under field conditions, *C. maculata* frequently consumes pollen (Forbes, 1881, 1883; Conrad, 1959; Putman, 1964; Solbreck, 1974; Benton and Crump, 1981; Lundgren et al., 2004), and lays more eggs in cropland during anthesis (Smith, 1971; Cottrell and Yeargan, 1998; Lundgren et al., 2004). Females consume 10 times more corn pollen than males under field conditions (Lundgren et al., 2005), and this is one of the few coccinellid species that can reproduce on a diet consisting solely of pollen (Lundgren and Wiedenmann, 2004). Clearly, pollinivory has contributed to the widespread abundance of this lady beetle across many habitats and geographic areas.

2.4. Other non-prey foods

A number of other non-prey foods are consumed by entomophagous coccinellids, but the importance of these foods to their diet is poorly understood. These foods include fruit, foliage, and fungus. Also, several gut dissections of field-collected lady beetles reveal that coccinellids frequently consume inorganic materials that are noteworthy (Putman, 1964; Anderson, 1982; Triltsch, 1999), but their role in the nutritional ecology of coccinellids is entirely unknown.

2.4.1. Fruit

Fruits are an especially good source of simple carbohydrates, and as such it is not uncommon to find lady beetles feeding at damaged fruit under field situations (Hodek and Honěk, 1996). Within the laboratory, providing fruits can improve the longevities of some coccinellids. For instance, raisins (accompanied by peach EFN) can help improve survival of *Stethorus punctillum* (Weise) in the laboratory (Putman, 1955), and *Anatis ocellata* (L.) consumed bananas as part of their laboratory diet (Kesten, 1969). The only empirical examination of fruit as food for coccinellids involves *Harmonia axyridis*, which is a secondary pest of several fruit crops in North America. This coccinellid frequently aggregates to fruit crops during the fall, where it damages grapes, apples, peaches, plums, pears, pumpkins and raspberries (Kovach, 2004; Koch and Galvan, 2008). The wine industry is particularly concerned by this insect, since pungent alkylmethoxy-pyrazines from even a single beetle can taint the flavor of wine (Kovach, 2004; Koch and Galvan, 2008). Fruit feeding by *H. axyridis* significantly improves their overwintering survival over unfed individuals (Galvan et al., 2009).

2.4.2. Foliage

Recent research shows (see also the report by Brassler, 1930) that coccinellids feed on plant foliage more often than previously believed, and thus are directly exposed to systemic insecticides and the insecticides produced by insect-tolerant GM crops. Moser et al. (2008) found that larvae (especially 4th instars) of *Coleomegilla maculata* and *Harmonia axyridis* fed on corn leaf tissue, even in the presence of water and aphids. Another laboratory assay found that *Harmonia axyridis* larvae (especially 1st and 4th instars) were adversely affected by soybean foliage possessing a soybean aphid (*Aphis glycines* Matsumura) resistance factor (the *Rag1* gene) (Lundgren et al.,

2009). In this study, host plant resistance-induced reductions in prey quality were not an influence on the experimental results, since lepidopteran eggs that had not ingested plant material were used as the prey item. These results suggest that there was a direct (and likely nutritional) effect of the soybean plant on the coccinellid.

2.4.3. Fungus

Fungus is a highly nutritious food source that is consumed by numerous entomophagous coccinellids (and fungal specialists; see Sutherland and Parrella, 2009). The most abundant constituent in most fungal tissues is water (85% by weight) (Chang and Miles, 2004). Spores are an exception to this trend, and typically contain minimal water. Although carbohydrates can comprise a substantial amount of the dry weight of fungus, most of this carbohydrate is in the form of structural polysaccharides (Mueller et al., 2001; Chang and Miles, 2004). Proteins are one of the most abundant constituents of fungal tissue, and fungi possess all of the amino acids essential to insect growth and development (Mueller et al., 2001; Chang and Miles, 2004). Lipids are often less abundant than carbohydrates and proteins, usually representing less than 10% of dry fungal weight (Harwood and Russell, 1984). Most of the sterols present in fungus are ergosterol, a C28 sterol that is usable by many insects. Vitamins and inorganic compounds are often quite abundant in fungi relative to other non-prey foods (Chang and Miles, 2004).

A range of entomophagous coccinellids consume fungal spores as an integral part of their diet, and many believe that mycophagy may have been associated with the early evolution of aphidophagy (lady beetles first consumed the sooty molds on honeydew before consuming the honeydew-producing sternorrhynchans; Sutherland and Parrella, 2009). Spores of at least 17 genera of fungus have been reported from the guts of coccinellids (Lundgren, 2009). Indeed, gut content analysis reveals that fungal spores are fed upon more frequently than prey by *Coccinella septempunctata* L., especially during the spring and fall (Tritsch, 1999). Another coccinellid, *Hippodamia convergens* Guérin-Méneville, consumes the spores of the plant pathogen, *Discula destructiva* Redlin, and is an important agent of its dispersal (Hed et al., 1999). Other species also use fungus to help build reserves before entering overwintering (Anderson, 1982; Ricci et al., 1983, 2005).

3. Nutritional suitability of non-prey foods for coccinellids

Lundgren (2009) statistically compared the nutritional contents (carbohydrates, protein, lipids, and calories) of hundreds of species of floral nectar, pollen, fungus, and arthropod prey. Pollen is quantitatively superior to prey in terms of energy (calories), protein, and carbohydrates, and is statistically similar to prey in lipid content. Floral nectar is deficient in lipids, proteins, and energy compared with prey, but has nearly three times the carbohydrate content. Fungus is statistically similar to prey in carbohydrate content, but is substantially lower in energy, protein, and lipid contents. Given the high nutritional content of some non-prey foods, it might be expected that coccinellids reared on non-prey foods such as pollen should have similar or superior fitness compared to those reared on prey. But considerable variability in the methods used to explore these questions experimentally, species-specific responses of coccinellids to prey and non-prey foods, and the nutritional properties of the non-prey foods themselves have challenged our ability to characterize trends in the nutritional suitability of non-prey foods for coccinellids.

Meta-analysis quantitatively and objectively examines cross-study trends in the literature to identify relationships that may be obscured or entirely overlooked by traditional literature reviews that either selectively report illustrative results of individual studies, or that use vote-counting approaches (presenting the proportion of studies that report a phenomenon) to characterize an ecological process (Bigger and Marvier, 1998; Osenberg et al., 1999; Rosenberg et al., 2000; Wolfenbarger et al., 2008). Here, meta-analysis was used to specifically address two questions:

- (1) Are prey or non-prey foods more suitable for coccinellids?
- (2) Are prey-only diets and mixed diets containing prey and non-prey foods equally suitable for coccinellids?

Two databases were created from the published literature to address these questions. The first compiled a series of 46 experimental comparisons from 14 published studies that made direct comparisons of a coccinellid species fed prey or non-prey diets (Table A.1). Ten coccinellid species and 11 prey species (Hemiptera: Sternorrhyncha [$n = 18$ studies], Lepidoptera: Phycitidae and Noctuidae [13], Coleoptera: Chrysomelidae and Curculionidae [8], Hymenoptera: Apidae [6]), were examined in this first database. The second database compiles 52 experimental comparisons from 16 published studies comparing prey-only to mixed diets on coccinellid performance and fitness (Table A.2). The second database includes 10 coccinellid species and nine prey species (Lepidoptera: Phycitidae and Noctuidae [$n = 27$ studies], Hemiptera: Sternorrhyncha [13], Coleoptera: Curculionidae [10], Hymenoptera: Apidae [2]).

The numerous life-history parameters measured in these studies were categorized as larval performance, adult performance, and reproduction. Some studies measured several aspects of these categories within a single experiment. To avoid non-independence within the database, only a single measurement was selected for each study for each category. For larval performance, larval developmental rate was prioritized over weight at eclosion. For adult




Fig. 1. The effects of prey versus non-prey foods (sugar and pollen analyzed simultaneously) on the larval and adult performance of coccinellid beetles. Effects are measured by Hedges' d (see text), with negative effect size indicating that the non-prey foods were inferior to the prey-only diet. Error bars represent 95% confidence intervals, and the arabic number associated with each bar represents the number of experimental comparisons analyzed.




Fig. 2. The effects of prey versus non-prey foods (sugar and pollen analyzed independently) on the larval and adult performance of coccinellid beetles. Effects are measured by Hedges' *d* (see text), with negative effect size indicating that the non-prey foods were inferior to the prey-only diet. Error bars represent 95% confidence intervals, and the arabic number associated with each bar represents the number of experimental comparisons analyzed.




Fig. 3. The effect of mixing prey-only diets with non-prey foods (sugar and pollen analyzed simultaneously) on coccinellid larvae and adults. Effects are measured by Hedges' *d* (see text), with negative effect size indicating that the mixed diets were inferior to the prey-only diet. Error bars denote 95% confidence intervals, arabic numbers associated with each bar indicate the number of experimental comparisons analyzed.

performance, adult longevity was prioritized over adult weight change. For reproduction, lifetime fecundity was compared preferentially over ovipositional or pre-ovipositional period. For the comparison of prey and non-prey diets, there are few reports where coccinellids are able to lay eggs on non-prey foods without prey (Hodek et al., 1978; Lundgren and Wiedenmann, 2004; Omkar, 2006; Berkvens et al., 2008), and the effects of prey versus non-prey foods on reproduction were not analyzed for this comparison. In studies that evaluated the effects of Bt and non-Bt pollen on coccinellids, the non-Bt hybrid was preferentially selected for analysis.

The meta-analyses used Hedges' *d* as its effect size estimator (Hedges and Olkin, 1985), with relative effect sizes assigned to each study based on the sample sizes, means and standard deviations of the two treatments compared. Contrasts between treatments were conducted such that a positive effect size represents a beneficial effect of the mixed or non-prey diet over the prey-only control diet. Comparisons were made using MetaWin 2.1, and mean \pm non-parametric bias-corrected bootstrap confidence intervals (representing 95% confidence limits) were calculated for each life-history parameter (Rosenberg et al., 2000). If the error intervals encompassed zero, the effect size was not considered to be significant. Small, medium, and large effect sizes were considered to be approximately 0.2, 0.4, and 0.6, respectively (Cohen, 1988). First, comparisons were made between prey-only diets and those involving non-prey foods as a whole. Then, the data was partitioned to determine whether pollen and sugar affected fitness of coccinellids similarly.

3.1. The relative suitability of prey and non-prey foods

Non-prey foods are clearly and substantially less suitable than prey for coccinellid larval and adult performance (Fig. 1). Indeed, sugar-fed larvae seldom complete development through the second stadium, and so only pollen-fed larvae were included in the analysis on larval performance (Figs. 1 and 2).




Fig. 4. The effect of mixing prey-only diets with sugar or pollen (non-prey foods analyzed independently) on coccinellid larvae and adults. Effects are measured by Hedges' *d* (see text), with negative effect size indicating that the non-prey foods were inferior to the prey-only diet. There was not a sufficient number of studies that compared prey with sugar in the larval performance for analysis. Error bars denote 95% confidence intervals, arabic numbers associated with each bar indicate the number of experiments analyzed. Asterisks indicate significant heterogeneity in the data ($P < 0.0001$).

What was surprising is that the longevity or weight change of sugar-fed coccinellids did not differ significantly from prey-fed beetles (Fig. 2). However, it should be noted that sugar is a poor food source for reproduction, and that many of the prey items tested in the literature are considered alternative foods. Although it has a greater breadth of nutrients, pollen was less effective in supporting adult performance than was prey. Two

non-exclusive possible explanations for these trends are that (1) pollen is limiting in unidentified nutrients critical to lady beetle performance, and that (2) pollen defenses or the types of nutrients in pollen render it unsuitable for lady beetles in the absence of prey.

3.2. Non-prey foods as components of mixed diets

There is a strong and positive effect of mixing prey-only diets with pollen on larval performance (Figs. 3 and 4). This suggests that pollen contains certain nutrients that are not present in the wide breadth of prey evaluated in published studies, and that the nutrition in pollen helps to improve some normally “essential” foods. Another trend in the literature is that mixing prey-only diets with non-prey foods does not uniformly change the fitness and performance of adult coccinellids (Fig. 3). However, the relative effects of pollen and sugar on this life stage at least partially explain the heterogeneity present in the database.

When additions of pollen and sugar to prey-only diets were analyzed separately, it was clear that coccinellids are affected very differently by these two non-prey foods. Adult performance and reproduction were strongly and positively affected when sugar was added to prey-only diets (Fig. 4). It should be added that the prey used in nearly all of these comparisons was not sternorrhynchan, although it was not necessarily of uniformly poor quality (Table A.2). The benefits of sugar in the diet may have stemmed from its nutrition or from possible phagostimulatory effects that spurred consumption of the prey. In contrast, adult performance and reproduction was not significantly affected by the addition of pollen to prey-only diets. There was substantial heterogeneity in each of the datasets presented in Figs. 1 and 2–4 except for the effects of sugar on reproduction.

4. Conclusions

Most, if not all, coccinellids rely on non-prey foods as part of their life history. Under field conditions, even the most stereotypical entomophagous coccinellids consume sugar, pollen, fungus, fruit and vegetation, often as an integral part of their diets. Non-prey foods serve as fuels for migration, support survival in the absence of prey, improve reproductive capacity, and increase survival through quiescent and overwintering periods. The importance of non-prey foods to coccinellids provides opportunities for pest managers to manipulate the abundance of and ecological services provided by coccinellids as biological control agents. For example, use of sugar sprays or habitat management to increase vegetational diversity often accompanies increases in biological control of key pests by coccinellid predators (Wade et al., 2008; Evans, 2009; Lundgren, 2009).

Meta-analyses of published literature on the suitability of non-prey foods suggest that coccinellid larvae have more stringent nutritional requirements than adults. Larvae perform poorly on non-prey foods alone, and while prey alone support

successful development, mixing prey and non-prey foods leads to faster development and greater weight gain in coccinellid larvae than when they are reared on prey alone. Pollen is in some ways nutritionally superior to insect prey, and entomologists are only beginning to understand what nutrients and phytochemicals limit pollen’s suitability as a sole food source, while enhancing the suitability of diets mixed with prey.

Carbohydrates are clearly an important dietary component for adult coccinellids. Sugar alone permits survival and reduces weight loss in adult coccinellids, and adding sugar to prey-only diets improves adult performance and increases reproduction substantially. Surprisingly, pollen has a very different effect on adult coccinellids, and generally reduces adult performance compared to prey-only diets when offered individually. Pollen as a component of mixed diets has no effect on reproduction and adult performance over prey-only diets. There was substantial heterogeneity in the effects on all tested interactions (except in the effects of sugar in mixed diets on coccinellid reproduction; Fig. 4), suggesting that not all coccinellids behave similarly to non-prey foods and that prey suitability differs widely among studies. Nevertheless, the significant and often dramatic positive effects of non-prey foods on these three life-history parameters clearly show that prey and non-prey foods are both parts of an optimal diet for many coccinellid larvae.

Finally, the clear importance of “alternative” foods to the optimal diets of coccinellids makes it important not to misinterpret the definitions of alternative and essential foods. While “essential” foods enable reproduction and development, these foods are not necessarily optimal for achieving maximum reproduction and developmental rates. The meta-analysis shows that the suitability of essential foods is usually improved when they are mixed with other foods. The predictability and reliability of biological control programs will be enhanced when we understand the nutritional constraints faced by omnivorous biological control agents like coccinellids, such that resources integral to their survival and success are available within cropland.

Acknowledgements

I would like to thank the editorial board of Biological Control for allowing Don Weber and me to organize this special issue, and Ted Evans, Ivo Hodek, and Don Weber for their comments on an earlier draft of this manuscript. Mention of a commercial or proprietary product does not constitute endorsement by the USDA.

Appendix Appendix. A

See Tables A.1 and A.2.

Table A.1 Database included in the meta-analysis comparing prey-only controls with non-prey diets (sugar and pollen) on two life-history categories of coccinellids. Effect size (Hedges' *d*) and *Var(d)* were calculated from means, standard deviations, and sample sizes published in the reference, or from direct communications with the author. A negative effect size indicates that the non-prey diet was inferior to the prey diet.

Coccinellid species	Prey species	Non-prey food	Parameter measured	Effect size (Hedges' <i>d</i>)	<i>Var(d)</i>	Reference
Larval performance						
<i>Adalia bipunctata</i> (L.)	<i>Ephesthia kuehniella</i> Zeller	Bee pollen mixture	Development period (d)	-8.8804	1.4088	De Clercq et al. (2005)
<i>Adalia bipunctata</i>	<i>Acyrtosiphon pisum</i> (Harris)	Rosaceae pollen	Development period (d)	-8.1803	3.4346	Hemptinne and Desprets (1986)
<i>Coleomegilla maculata</i> (Pollen 2)	<i>Rhopalosiphum maidis</i> (Fitch)	Corn pollen	Development period (d)	-1.6914	0.1165	Lundgren and Wiedenmann (2004)
<i>Coleomegilla maculata</i> (Pollen 3)	<i>Rhopalosiphum maidis</i>	Corn pollen	Development period (d)	-1.8491	0.1225	Lundgren and Wiedenmann (2004)
<i>Coleomegilla maculata</i> (Pollen 4)	<i>Rhopalosiphum maidis</i>	Corn pollen	Development period (d)	-1.6506	0.113	Lundgren and Wiedenmann (2004)
<i>Coleomegilla maculata</i>	<i>Rhopalosiphum maidis</i>	Corn pollen	Development period (d)	-3.5384	0.855	Smith (1965)
<i>Coleomegilla maculata</i>	<i>Myzus persicae</i> (Sulzer)	Corn pollen	Development period (d)	-1.2977	0.1729	Hazzard and Ferro (1991)
<i>Coleomegilla maculata</i>	<i>Leptinotarsa decemlineata</i> (Say)	Corn pollen	Development period (d)	2.3316	0.2711	Hazzard and Ferro (1991)
<i>Coleomegilla maculata</i>	<i>Schizaphis graminum</i> (Rondani)	Bee pollen	Development period (d)	-9.4452	0.4192	Michaud and Jyoti (2008)
<i>Coleomegilla maculata</i>	<i>Ephesthia kuehniella</i>	Bee pollen	Development period (d)	-8.3675	0.3485	Michaud and Jyoti (2008)
<i>Harmonia axyridis</i> (laboratory females)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	Development period (d)	-4.832	0.3148	Berkvens et al. (2008)
<i>Harmonia axyridis</i> (red females)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	Development period (d)	-5.3429	0.4343	Berkvens et al. (2008)
<i>Harmonia axyridis</i> (black females)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	Development period (d)	-3.7606	0.2484	Berkvens et al. (2008)
<i>Harmonia axyridis</i> (laboratory males)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	Development period (d)	-5.2423	0.5326	Berkvens et al. (2008)
<i>Harmonia axyridis</i> (red males)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	Development period (d)	-8.3538	1.3322	Berkvens et al. (2008)
<i>Harmonia axyridis</i> (black males)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	Development period (d)	-6.2844	0.7042	Berkvens et al. (2008)
<i>Harmonia axyridis</i>	<i>Myzus persicae</i>	Corn pollen	Development period (d)	-9.4045	5.0473	Hukusima and Itoh (1976)
<i>Harmonia axyridis</i>	<i>Myzus persicae</i>	Corn pollen + honey	Development period (d)	-5.1101	1.0041	Hukusima and Itoh (1976)
<i>Harmonia axyridis</i>	<i>Apis mellifera</i> L.	Corn pollen	Development period (d)	-5.5991	1.3838	Hukusima and Itoh (1976)
<i>Harmonia axyridis</i>	<i>Apis mellifera</i>	Corn pollen + honey	Development period (d)	-3.6083	0.4203	Hukusima and Itoh (1976)
<i>Harmonia axyridis</i>	<i>Rhopalosiphum maidis</i>	Corn pollen	Development period (d)	-0.9644	0.2232	Hukusima and Itoh (1976)
<i>Micraspis discolor</i> (Fabricius)	<i>Aphis gossypii</i> Glover F1	Corn pollen	Development period (d)	1.1139	0.165	Omkar (2006)
<i>Micraspis lineata</i> (Thunberg)	<i>Aphis gossypii</i> F2	Bee pollen	Development period (d)	-3.1769	0.165	Anderson and Hales (1983)
<i>Micraspis lineata</i>	<i>Acyrtosiphon pisum</i>	Bee pollen	Development period (d)	0.3489	0.3489	Anderson and Hales (1983)
<i>Micraspis lineata</i>	<i>Apis mellifera</i>	Bee pollen	Development period (d)	0.2692	0.2692	Anderson and Hales (1983)
<i>Propylea japonica</i> (Thunberg)	<i>Myzus persicae</i>	Rye pollen	Development period (d)	-0.1615	0.1433	Anderson and Hales (1983)
<i>Propylea japonica</i>	<i>Myzus persicae</i>	Corn pollen + honey	Development period (d)	-5.3752	1.2546	Hukusima and Itoh (1976)
<i>Propylea japonica</i>	<i>Apis mellifera</i>	Rye pollen	Development period (d)	-4.5555	0.7569	Hukusima and Itoh (1976)
<i>Propylea japonica</i>	<i>Apis mellifera</i>	Corn pollen + honey	Development period (d)	-4.2874	0.9454	Hukusima and Itoh (1976)
<i>Propylea japonica</i>	<i>Apis mellifera</i>	Corn pollen + honey	Development period (d)	-4.2135	0.6819	Hukusima and Itoh (1976)
Adult performance						
<i>Adalia bipunctata</i> (Exp. III)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	Weight gain (mg)	-4.3077	0.5474	De Clercq et al. (2005)
<i>Coccinella septempunctata</i> (Exp. IV, females)	<i>Hypera postica</i> Gyllenhal	Sugar	Weight change (mg)	-0.3539	0.1354	Richards and Evans (1998)
<i>Coccinella septempunctata</i> (Exp. IV, males)	<i>Hypera postica</i>	Sugar	Weight change (mg)	-0.9602	0.1394	Richards and Evans (1998)
<i>Coccinella septempunctata</i> (Exp. VI, females)	<i>Hypera postica</i>	Sugar	Weight change (mg)	-0.1727	0.1673	Richards and Evans (1998)
<i>Coccinella septempunctata</i> (Exp. VI, males)	<i>Hypera postica</i>	Sugar	Weight change (mg)	-1.2574	0.2994	Richards and Evans (1998)
<i>Coccinella transversalis</i>	<i>Myzus persicae</i>	Sugar	Weight change (mg)	-2.1611	0.5279	Evans (2000)
<i>Coccinella transversalis</i>	<i>Helicoverpa armigera</i> (Hübner)	Sugar	Weight change (mg)	0.2004	0.335	Evans (2000)
<i>Coccinella transversoguttata</i> Falderman (Exp. IV, females)	<i>Hypera postica</i>	Sugar	Weight change (mg)	-1.4877	0.2574	Richards and Evans (1998)
<i>Coccinella transversoguttata</i> (Exp. IV, males)	<i>Hypera postica</i>	Sugar	Weight change (mg)	-2.1065	0.2966	Richards and Evans (1998)
<i>Harmonia axyridis</i> (laboratory population)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	Longevity (d)	-1.3202	0.107	Richards and Evans (1998)
<i>Harmonia axyridis</i> (red population)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	Longevity (d)	-0.6189	0.1117	Berkvens et al. (2008)
<i>Harmonia axyridis</i> (black population)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	Longevity (d)	-1.8719	0.141	Berkvens et al. (2008)
<i>Harmonia axyridis</i>	<i>Hypera postica</i>	Sugar	Weight change (mg)	-2.0175	0.3772	Evans and Gunther (2005)
<i>Harmonia axyridis</i>	<i>Acyrtosiphon pisum</i>	Sugar	Weight change (mg)	-5.157	1.0811	Evans and Gunther (2005)
<i>Rodolia cardinalis</i> (Mulsant)	<i>Icerya purchasi</i> Maskell	Sucrose	Longevity (d)	0.5884	0.1043	Matsuka et al. (1982)
<i>Rodolia cardinalis</i>	<i>Apis mellifera</i>	Sucrose	Longevity (d)	2.703	0.1913	Matsuka et al. (1982)

Table A.2 Database included in the meta-analysis comparing prey-only controls with mixed diets (prey + non-prey foods) on three life-history categories of coccinellids. Effect size (Hedges' *d*) and Var(*d*) were calculated from means, standard deviations, and sample sizes published in the reference, or from direct communications with the author. A negative effect size indicates that the mixed diet was inferior to the prey-only diet.

Coccinellid species	Prey species	Non-prey food	Parameter measured	Effect size (Hedges' <i>d</i>)	Var(<i>d</i>)	Reference
Larval performance						
<i>Adalia bipunctata</i> Exp. III	<i>Ephesthia kuehniella</i>	Bee pollen mixture	Development period (d)	1.0121	0.0728	De Clercq et al. (2005)
<i>Adalia bipunctata</i> Exp. II	<i>Ephesthia kuehniella</i>	Dry bee pollen mixture	Development period (d)	0	0.0774	De Clercq et al. (2005)
<i>Adalia bipunctata</i> Exp. II	<i>Ephesthia kuehniella</i>	Frozen bee pollen mixture	Development period (d)	-0.305	0.0728	De Clercq et al. (2005)
<i>Adalia bipunctata</i>	<i>Acyrtosiphon pisum</i>	Rosaceae pollen	Development period (d)	1.219	0.3174	Hemphime and Desprets (1986)
<i>Coleomegilla maculata</i>	<i>Rhopalosiphum maidis</i>	Corn pollen	Development period (d)	3.9205	0.2921	Smith (1965)
<i>Coleomegilla maculata</i>	<i>Rhopalosiphum maidis</i>	Corn pollen	Development period (d)	1.1096	0.2308	Omkar (2006)
<i>Coleomegilla maculata</i>	<i>Myzus persicae</i>	Rice pollen	Development period (d)	-1.1645	0.0557	Bai et al. (2005)
<i>Harmonia axyridis</i> (laboratory females)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	Development period (d)	1.4913	0.0787	Berkvens et al. (2008)
<i>Harmonia axyridis</i> (red females)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	Development period (d)	0.0202	0.067	Berkvens et al. (2008)
<i>Harmonia axyridis</i> (black females)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	Development period (d)	3.1458	0.1494	Berkvens et al. (2008)
<i>Harmonia axyridis</i> (laboratory males)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	Development period (d)	1.1658	0.0936	Berkvens et al. (2008)
<i>Harmonia axyridis</i> (red males)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	Development period (d)	-0.4368	0.0897	Berkvens et al. (2008)
<i>Harmonia axyridis</i> (black males)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	Development period (d)	2.8001	0.1553	Berkvens et al. (2008)
<i>Harmonia axyridis</i>	<i>Apis mellifera</i>	Brewer's yeast (50% of diet) + sucrose	Development period (d)	1.274	0.1203	Nijima et al. (1997)
<i>Micraspis discolor</i>	<i>Acyrtosiphon pisum</i>	Corn pollen	Development period (d)	1.7389	0.0919	Picher et al. (1997)
<i>Propylea japonica</i>	<i>Schizaphis graminum</i>	Corn pollen	Development period (d)	1.2066	0.0249	Ahmad et al. (2006)
Adult performance						
<i>Adalia bipunctata</i> (Exp. III)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	Weight gain (mg)	3.2151	0.2292	De Clercq et al. (2005)
<i>Adalia bipunctata</i> (Exp. II)	<i>Ephesthia kuehniella</i>	Dry bee pollen mixture	Weight gain (mg)	0.3509	0.0655	De Clercq et al. (2005)
<i>Adalia bipunctata</i> (Exp. II)	<i>Ephesthia kuehniella</i>	Frozen bee pollen mixture	Weight gain (mg)	0.213	0.0778	De Clercq et al. (2005)
<i>Coccinella septempunctata</i> (Exp. IV, females)	<i>Hypera postica</i>	Sucrose	Weight change (mg)	0.9217	0.1429	Richards and Evans (1998)
<i>Coccinella septempunctata</i> (Exp. IV, males)	<i>Hypera postica</i>	Sucrose	Weight change (mg)	0.1614	0.1217	Richards and Evans (1998)
<i>Coccinella septempunctata</i> (Exp. VI, females)	<i>Hypera postica</i>	Sucrose	Weight change (mg)	1.1311	0.2403	Richards and Evans (1998)
<i>Coccinella septempunctata</i> (Exp. VI, males)	<i>Hypera postica</i>	Sucrose	Weight change (mg)	0.2051	0.217	Richards and Evans (1998)
<i>Coccinella transversalis</i>	<i>Helicoverpa armigera</i>	Sucrose	Weight change (mg)	0.044	0.3334	Evans (2000)
<i>Coccinella transversalis</i>	<i>Helicoverpa armigera</i>	Aphid honeydew	Weight change (mg)	0.3842	0.3395	Evans (2000)
<i>Coccinella transversoguttata</i> (females)	<i>Hypera postica</i>	Sucrose	Weight change (mg)	-0.0426	0.2223	Richards and Evans (1998)
<i>Coccinella transversoguttata</i> (males)	<i>Hypera postica</i>	Sucrose	Weight change (mg)	-0.3561	0.2032	Richards and Evans (1998)
<i>Coleomegilla maculata</i>	<i>Schizaphis graminum</i>	Corn pollen	Longevity (d)	-0.182	0.0283	Ahmad et al. (2006)
<i>Harmonia axyridis</i> (laboratory population)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	Longevity (d)	-1.0745	0.0776	Berkvens et al. (2008)
<i>Harmonia axyridis</i> (red population)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	Longevity (d)	-0.362	0.0713	Berkvens et al. (2008)
<i>Harmonia axyridis</i> (black population)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	Longevity (d)	-0.5994	0.0733	Berkvens et al. (2008)
<i>Harmonia axyridis</i>	<i>Hypera postica</i>	Sucrose	Weight change (mg)	-0.9146	0.2761	Evans and Gunther (2005)
<i>Hippodamia convergens</i>	<i>Schizaphis graminum</i>	Bee pollen mixture	Longevity (d)	-0.1092	0.0589	Michaud and Qureshi (2006)
<i>Rodolia cardinalis</i>	<i>Icerya purchasi</i>	Sucrose	Longevity (d)	3.4656	0.2501	Matsuka et al. (1982)
<i>Rodolia cardinalis</i>	<i>Apis mellifera</i>	Sucrose	Longevity (d)	3.2151	0.2292	Matsuka et al. (1982)
Reproduction						
<i>Adalia bipunctata</i> (Exp. III)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	# eggs	1.198	0.2286	De Clercq et al. (2005)
<i>Adalia bipunctata</i> (Exp. II)	<i>Ephesthia kuehniella</i>	Dry bee pollen mixture	# eggs	0.3919	0.238	De Clercq et al. (2005)
<i>Adalia bipunctata</i> (Exp. II)	<i>Ephesthia kuehniella</i>	Frozen bee pollen mixture	# eggs	0.6981	0.232	De Clercq et al. (2005)
<i>Adalia bipunctata</i>	<i>Acyrtosiphon pisum</i>	Rosaceae pollen	# eggs	-0.2499	0.2699	Hemphime and Desprets (1986)
<i>Coccinella septempunctata</i>	<i>Hypera postica</i>	Sucrose	# eggs	1.2846	0.1558	Richards and Evans (1998)
<i>Coccinella transversalis</i>	<i>Helicoverpa armigera</i>	Sucrose	# eggs	0.9332	0.3696	Evans (2000)
<i>Coccinella transversalis</i>	<i>Helicoverpa armigera</i>	Aphid honeydew	# eggs	0.7008	0.3538	Evans (2000)
<i>Coccinella transversoguttata</i>	<i>Hypera postica</i>	Sucrose	# eggs	1.231	0.2643	Richards and Evans (1998)
<i>Coleomegilla maculata</i>	<i>Ephesthia kuehniella</i>	Corn pollen	# eggs	0.0336	0.1181	Richards and Evans (1998)
<i>Coleomegilla maculata</i>	<i>Ephesthia kuehniella</i>	Sorghum pollen	# eggs	0.2196	0.1151	Michaud and Grant (2005)
<i>Harmonia axyridis</i> (laboratory population)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	# eggs	-0.2298	0.076	Berkvens et al. (2008)
<i>Harmonia axyridis</i> (red population)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	# eggs	0.1481	0.0819	Berkvens et al. (2008)
<i>Harmonia axyridis</i> (black population)	<i>Ephesthia kuehniella</i>	Bee pollen mixture	# eggs	-0.0622	0.0917	Berkvens et al. (2008)
<i>Harmonia axyridis</i>	<i>Hypera postica</i>	Sucrose	# eggs	0.9881	0.2805	Evans and Gunther (2005)
<i>Hippodamia convergens</i>	<i>Schizaphis graminum</i>	Bee pollen mixture	# eggs	-2.4859	0.0821	Michaud and Qureshi (2006)
<i>Micraspis discolor</i>	<i>Rhopalosiphum maidis</i>	Corn pollen	# eggs	2.4475	0.3498	Omkar (2006)
<i>Propylea japonica</i>	<i>Myzus persicae</i>	Rice pollen	# eggs	0.7462	0.1535	Bai et al. (2005)

References

- Adler, L.S., 2000. The ecological significance of toxic nectar. *Oikos* 91, 409–420.
- Adler, L.S., Irwin, R.E., 2005. Ecological costs and benefits of defenses in nectar. *Ecology* 86, 2968–2978.
- Ahmad, A., Wilde, G.E., Whitworth, R.J., Zolnerowich, G., 2006. Effect of corn hybrids expressing the coleopteran-specific Cry3Bb1 protein for corn rootworm control on aboveground insect predators. *Journal of Economic Entomology* 99, 1085–1095.
- Anderson, J.M.E., 1981. Seasonal field analyses of fat content, live weight, dry weight and water content of the aphidophagous *Scymnodes lividigaster* (Mulsant) and mycophagous *Leptothea galbula* (Mulsant) (Coleoptera: Coccinellidae). *Australian Journal of Zoology* 29, 679–689.
- Anderson, J.M.E., 1982. Seasonal habitat utilization and food of the ladybirds *Scymnodes lividigaster* (Mulsant) and *Leptothea galbula* (Mulsant) (Coleoptera: Coccinellidae). *Australian Journal of Zoology* 30, 59–70.
- Anderson, J.M.E., Hales, D.F., 1983. *Micraspis lineata* (Thunberg) (Coleoptera: Coccinellidae) – seasonality and food. *General and Applied Entomology* 15, 47–52.
- Ashford, D.A., Smith, W.A., Douglas, A.E., 2000. Living on a high sugar diet: the fate of sucrose ingested by a phloem-feeding insect, the pea aphid *Acyrtosiphon pisum*. *Journal of Insect Physiology* 46, 335–341.
- Bai, Y.Y., Jiang, M.X., Cheng, J.A., 2005. Effects of transgenic cry1Ab rice pollen on fitness of *Propylea japonica* (Thunberg). *Journal of Pest Science* 78, 123–128.
- Baker, H.G., 1975. Sugar concentrations in nectars from hummingbird flowers. *Biotropica* 7, 37–41.
- Baker, H.G., Baker, I., 1977. Intraspecific constancy of floral nectar amino acid complements. *Botanical Gazette* 138, 183–191.
- Baker, H.G., Baker, I., 1978. Ants and flowers. *Biotropica* 10, 80.
- Baker, H.G., Baker, I., 1979. Starch in angiosperm pollen grains and its evolutionary significance. *American Journal of Botany* 66, 591–600.
- Baker, H.G., Baker, I., 1982. Starchy and starchless pollen in the Onagraceae. *Annals of the Missouri Botanical Garden* 69, 748–754.
- Baker, H.G., Baker, I., 1983. A brief historical review of the chemistry of floral nectar. In: Bentley, B.L., Elias, T.S. (Eds.), *A Brief Historical Review of The Chemistry of Floral Nectar*. Columbia University Press, New York, pp. 126–152.
- Baker, H.G., Opler, P.A., Baker, I., 1978. A comparison of the amino acid complements of floral and extrafloral nectars. *Botanical Gazette* 139, 322–332.
- Banks, C.J., 1957. The behaviour of individual coccinellid larvae on plants. *Animal Behaviour* 5, 12–24.
- Barbier, M., 1970. Chemistry and biochemistry of pollens. *Progress in Phytochemistry* 2, 1–33.
- Bentley, B.L., 1977. Extrafloral nectaries and protection by pugnacious bodyguards. *Annual Review of Ecology and Systematics* 8, 407–427.
- Benton, A.H., Crump, A.J., 1981. Observations on the spring and summer behavior of the 12-spotted ladybird beetle, *Coleomegilla maculata* (DeGeer) (Coleoptera: Coccinellidae). *New York Entomological Society* 89, 102–108.
- Berkvens, N., Bonte, J., Berkvens, D., Deforce, K., Tirry, L., De Clercq, P., 2008. Pollen as an alternative food for *Harmonia axyridis*. *BioControl* 53, 201–210.
- Bigger, D.S., Marvier, M., 1998. How different would a world without herbivory be? A search for generality in ecology. *Integrative Biology* 1, 60–67.
- Brassler, K., 1930. Ist *Coccinella septempunctata* L. wirklich nur Blattlausfresser? *Zeitschrift für Pflanzenkrankheit, Pflanzenpathologie, und Pflanzenschutz* 40, 511–513.
- Bristow, C.M., 1991. Are ant-aphid associations a tritrophic interaction? *Oleander aphids and Argentine ants*. *Oecologia* 87, 514–521.
- Bugg, R.L., 1987. Observations on insects associated with a nectar-bearing Chilean tree, *Quillaja saponaria* Molina (Rosaceae). *Pan-Pacific Entomologist* 63, 60–64.
- Caldwell, D.L., Gerhardt, K.O., 1986. Chemical analysis of peach extrafloral nectary exudate. *Phytochemistry* 25, 411–413.
- Carter, C., Shafir, S., Yehonatan, L., Palmer, R.G., Thornburg, R., 2006. A novel role for proline in plant floral nectars. *Naturwissenschaften* 93, 72–79.
- Carter, M.C., Dixon, F.G., 1984. Honeydew: an arrestant stimulus for coccinellids. *Ecological Entomology* 9, 383–387.
- Chang, S.-T., Miles, P.G., 2004. *Mushrooms: Cultivation, Nutritional Value, Medicinal Effect, and Environmental Impact*. CRC Press, Boca Raton, FL.
- Choh, Y., Kugimiya, S., Takabayashi, J., 2006. Induced production of extra-floral nectar in intact lima bean plants in response to volatiles from spider mite-infested conspecific plants as a possible indirect defense against spider mites. *Oecologia* 147, 455–460.
- Cohen, J., 1988. *Statistical Power Analysis for the Behavioral Sciences*, second ed. Lawrence Erlbaum Associates.
- Conrad, M.S., 1959. The spotted lady beetle, *Coleomegilla maculata* (De Geer), as a predator of European corn borer eggs. *Journal of Economic Entomology* 52, 843–847.
- Costa, H.S., Toscano, N.C., Hendrix, D.L., Henneberry, T.J., 1999. Patterns of honeydew droplet production by nymphal stages of *Bemisia argentifolii* (Homoptera: Aleyrodidae) and relative composition of honeydew sugars. *Journal of Entomological Science* 34, 305–313.
- Cottrell, T.E., Yeargan, K.V., 1998. Effect of pollen on *Coleomegilla maculata* (Coleoptera: Coccinellidae) population density, predation, and cannibalism in sweet corn. *Environmental Entomology* 27, 1402–1410.
- Cruden, R.W., Hermann, S.M., Peterson, S., 1983. Patterns of nectar production and plant-pollinator coevolution. In: Bentley, B.L., Elias, T.S. (Eds.), *Patterns of Nectar Production and Plant-Pollinator Coevolution*. Columbia University Press, New York, pp. 80–125.
- De Clercq, P., Bonte, M., Van Speybroeck, K., Bolckmans, K., Deforce, K., 2005. Development and reproduction of *Adalia bipunctulata* (Coleoptera: Coccinellidae) on eggs of *Ephestia kuehniella* (Lepidoptera: Phycitidae) and pollen. *Pest Management Science* 61, 1129–1132.
- DeVries, P.J., Baker, I., 1989. Butterfly exploitation of an ant-plant mutualism: adding insult to herbivory. *Journal of the New York Entomological Society* 97, 332–340.
- Downes Jr., W.L., Dahlem, G.A., 1987. Keys to the evolution of Diptera: role of Homoptera. *Environmental Entomology* 16, 847–854.
- Dreyer, B.S., Neuenschwander, P., Baumgartner, J., Dorn, S., 1997. Trophic influences on survival, development and reproduction of *Hyperaspis notata* (Col., Coccinellidae). *Journal of Applied Entomology* 121, 249–256.
- Elias, T.S., Gelband, H., 1975. Nectar: its production and functions in trumpet creeper. *Science* 189, 289–291.
- Erhardt, A., Baker, I., 1990. Pollen amino acids – an additional diet for a nectar feeding butterfly? *Plant Systematics and Evolution* 169, 111–121.
- Evans, E.W., 2000. Egg production in response to combined alternative foods by the predator *Coccinella transversalis*. *Entomologia Experimentalis et Applicata* 94, 141–147.
- Evans, E.W., 2009. Lady beetles as predators of insects other than Hemiptera. *Biological Control* 51, 255–267.
- Evans, E.W., Gunther, D.I., 2005. The link between food and reproduction in aphidophagous predators: a case study with *Harmonia axyridis* (Coleoptera: Coccinellidae). *European Journal of Entomology* 102, 423–430.
- Ewart, W.H., Metcalf, R.L., 1956. Preliminary studies of sugars and amino acids in the honeydew of five species of coccids feeding on citrus in California. *Annals of the Entomological Society of America* 49, 441–447.
- Ewing, H.E., 1913. Notes on oregon coccinellidae. *Journal of Economic Entomology* 6, 404–407.
- Forbes, S.A., 1881. Notes on insectivorous Coleoptera. *Bulletin of the Illinois State Laboratory of Natural History* 1, 153–160.
- Forbes, S.A., 1883. The food relations of the Carabidae and Coccinellidae. *Bulletin of the Illinois State Laboratory of Natural History* 3, 33–64.
- Galletto, L., Bernardello, G., 2004. Floral nectaries, nectar production dynamics and chemical composition in six *Ipomoea* species (Convolvulaceae) in relation to pollinators. *Annals of Botany* 94, 269–280.
- Galvan, T.L., Koch, R.L., Hutchison, W.D., 2009. Impact of fruit feeding on overwintering survival of the multicolored Asian lady beetle, and the ability of this insect and paper wasps to injure wine grape berries. *Entomologia Experimentalis et Applicata*. doi:10.1111/j.1570-7458.2008.00731.x.
- Gardener, M.C., Gillman, M.P., 2002. The taste of nectar – a neglected area of pollination ecology. *Oikos* 98, 552–557.
- Geyer, J.W.C., 1947. A study of the biology and ecology of *Exochomus flavipes* Thunb. (Coccinellidae, Coleoptera). *Journal of the Entomological Society of South Africa* 9, 219–234.
- Giorgi, J.A., Vandenberg, N.J., McHugh, J.V., Forrester, J.A., Ślipiński, S.A., Miller, K.B., Shapiro, I.R., Whiting, 2009. The evolution of food preferences in Coccinellidae. *Biological Control* 51, 215–231.
- Gottsberger, G., Schrauwen, J., Linskins, H.F., 1984. Amino acids and sugars in nectar, and their putative evolutionary significance. *Plant Systematics and Evolution* 145, 55–77.
- Gray, R.A., 1952. Composition of honeydew excreted by pineapple mealybugs. *Science* 115, 129–133.
- Guerrant Jr., E.O., Fiedler, P.L., 1981. Flower defenses against nectar-pilferage by ants. *Biotropica* 13, 25–33.
- Hagen, K.S., 1962. Biology and ecology of predaceous Coccinellidae. *Annual Review of Entomology* 7, 289–326.
- Hagen, K.S., 1986. Ecosystem analysis: plant cultivars (HPR), entomophagous species and food supplements. In: Boethel, D.J., Eikenbary, R.D. (Eds.), *Ecosystem Analysis: Plant Cultivars (HPR), Entomophagous Species and Food Supplements*. Ellis Horwood Publishers, Chichester, UK, pp. 151–197.
- Han, B., Chen, Z., 2002. Behavioral and electrophysiological responses of natural enemies to synomones from tea shoots and kairomones from tea aphids, *Toxoptera aurantii*. *Journal of Chemical Ecology* 28, 2203–2219.
- Hanney, B.W., Elmore, C.D., 1974. Amino acid composition of cotton nectar. *Journal of Agricultural and Food Chemistry* 22, 476–478.
- Harwood, J.L., Russell, N.J., 1984. *Lipids in Plants and Microbes*. George Allen and Unwin, London, UK.
- Hazzard, R.V., Ferro, D.N., 1991. Feeding responses of adult *Coleomegilla maculata* (Coleoptera: Coccinellidae) to eggs of Colorado potato beetle (Coleoptera: Chrysomelidae) and green peach aphids (Homoptera: Aphididae). *Environmental Entomology* 20, 644–651.
- Hed, B.E., Windham, M.T., Grant, J.F., 1999. Survival of conidia of *Discula destructiva* in frass of the convergent lady beetle. *Plant Disease* 83, 806–809.
- Hedges, L.V., Olkin, I., 1985. *Statistical Methods for Meta-Analysis*. Academic Press, New York, NY.
- Heil, M., 2004. Induction of two indirect defences benefits lima bean (*Phaseolus lunatus*, Fabaceae) in nature. *Journal of Ecology* 92, 527–536.
- Heil, M., Fiala, B., Baumann, B., Linsenmair, K.E., 2000. Temporal, spatial and biotic variations in extrafloral nectar secretion by *Macaranga tanarius*. *Functional Ecology* 14, 749–757.
- Heil, M., Koch, T., Hilpert, A., Fiala, B., Boland, W., Linsenmair, K.E., 2001. Extrafloral nectar production of the ant-associated plant, *Macaranga tanarius*, is an

- induced, indirect, defensive response elicited by jasmonic acid. *Proceedings of the National Academy of Sciences* 98, 1083–1088.
- Hemptinne, J.L., Desprets, A., 1986. Pollen as Spring Food for *Adalia bipunctata*. In: Hodek, I. (Ed.), *Ecology of Aphidophaga*. Prague & Dr W. Junk, Dordrecht, The Netherlands, pp. 29–35.
- Hemptinne, J.L., Naisse, J., 1987. Ecology of the reproductive activity of *Adalia bipunctata* L. (Col., Coccinellidae). *Mededelingen Faculteit Landbouwwetenschappen Rijksuniversiteit Gent* 52, 225–233.
- Hodek, I., Honěk, A., 1996. *Ecology of Coccinellidae*. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Hodek, I., Honěk, A., 2009. Scale insects, psyllids, and whiteflies (Hemiptera: Sternorrhyncha) – prey of ladybirds. *Biological Control* 51, 232–243.
- Hodek, I., Růžička, Z., Hodková, M., 1978. Pollinivorie et aphidophagie chez *Coleomegilla maculata* (DeGeer). *Annales de Zoologie Ecologie Animale* 10, 453–459.
- Hoheisel, G.-A., Fleischer, S.J., 2007. Coccinellids, aphids, and pollen in diversified vegetable fields with transgenic and isoline cultivars. *Journal of Insect Science* 7, 61.
- Hukusima, S., Itoh, K., 1976. Pollen and fungus as food for some coccinellid beetles. *Research Bulletin of the Faculty of Agriculture at Gifu University* 39, 31–37.
- Ibrahim, M.M., 1955. Studies on *Coccinella undecimpunctata aegyptiaca* Reiche. 2. Biology and life-history. *Bulletin of the Entomological Society of Egypt* 39, 395–423.
- Jakubská, A., Przado, D., Steininger, M., Aniol-Kwiatkowska, J., Kadej, M., 2005. Why do pollinators become “sluggish”? Nectar chemical constituents from *Epipactis helleborine* (L.) Crantz (Orchidaceae). *Applied Ecology and Environmental Research* 3, 29–38.
- Keeler, K.H., 1977. The extrafloral nectaries of *Ipomoea carnea* (Convolvulaceae). *American Journal of Botany* 64, 1182–1188.
- Keeler, K.H., 1978. Insects feeding at extrafloral nectaries of *Ipomoea carnea* (Convolvulaceae). *Entomological News* 89, 163–168.
- Kessler, D., Gase, K., Baldwin, I.T., 2008. Field experiments with transformed plants reveal sense of floral scents. *Science* 321, 1200–1202.
- Kesten, U., 1969. Zur Morphologie und Biologie von *Anatis ocellata* (L.) (Coleoptera, Coccinellidae). *Zeitschrift für Angewandte Entomologie* 63, 412–455.
- Koch, R.L., Galvan, T.L., 2008. Bad side of a good beetle: the North American experience with *Harmonia axyridis*. *BiolControl* 53, 23–35.
- Kost, C., Heil, M., 2006. Herbivore-induced plant volatiles induce an indirect defence in neighbouring plants. *Journal of Ecology* 94, 619–628.
- Kovach, J., 2004. Impact of multicolored Asian lady beetles as a pest of fruit and people. *American Entomologist* 50, 159–161.
- Lamb, K.P., 1959. Composition of the honeydew of the aphid *Brevicoryne brassicae* (L.) feeding on swedes (*Brassica napobrassica* DC.). *Journal of Insect Physiology* 3, 1–13.
- Lundgren, J.G., 2009. *Relationships of Natural Enemies and Non-prey Foods*. Springer International, Dordrecht, The Netherlands.
- Lundgren, J.G., Hesler, L.S., Tilmon, K.M., Dashiell, K.E., Scott, R., 2009. Direct effects of soybean varietal selection and *Aphis glycines*-resistant soybeans on natural enemies. *Arthropod-Plant Interactions* 3, 9–16.
- Lundgren, J.G., Huber, A., Wiedenmann, R.N., 2005. Quantification of consumption of corn pollen by the predator *Coleomegilla maculata* (Coleoptera: Coccinellidae) during anthesis in an Illinois cornfield. *Agricultural and Forest Entomology* 7, 53–60.
- Lundgren, J.G., Razzak, A.A., Wiedenmann, R.N., 2004. Population responses and food consumption by predators *Coleomegilla maculata* and *Harmonia axyridis* (Coleoptera: Coccinellidae) during anthesis in an Illinois cornfield. *Environmental Entomology* 33, 958–963.
- Lundgren, J.G., Wiedenmann, R.N., 2004. Nutritional suitability of corn pollen for the predator *Coleomegilla maculata* (Coleoptera: Coccinellidae). *Journal of Insect Physiology* 50, 567–575.
- Majerus, M.E.N., Sloggett, J.J., Godeau, J.-F., Hemptinne, J.-L., 2007. Interactions between ants and aphidophagous and coccidophagous ladybirds. *Population Ecology* 49, 15–27.
- Matsuka, M., Watanabe, M., Nijima, K., 1982. Longevity and oviposition of *Vedalia* beetles on artificial diets. *Environmental Entomology* 11, 816–819.
- Michaud, J.P., 2000. Development and reproduction of ladybeetles (Coleoptera: Coccinellidae) on the citrus aphids *Aphis spiraecola* Patch and *Toxoptera citricida* (Kirkaldy) (Homoptera: Aphididae). *Biological Control* 18, 287–297.
- Michaud, J.P., Grant, A.K., 2005. Suitability of pollen sources for the development and reproduction of *Coleomegilla maculata* (Coleoptera: Coccinellidae) under simulated drought conditions. *Biological Control* 32, 363–370.
- Michaud, J.P., Jyoti, J.L., 2008. Dietary complementation across life stages in the polyphagous lady beetle *Coleomegilla maculata*. *Entomologia Experimentalis et Applicata* 126, 40–45.
- Michaud, J.P., Qureshi, J.A., 2006. Reproductive diapause in *Hippodamia convergens* (Coleoptera: Coccinellidae) and its life history consequences. *Biological Control* 39, 193–200.
- Mittler, T.E., 1958. Studies on the feeding and nutrition of *Tuberolachnus salignus* (Gmelin) (Homoptera: Aphididae). II. The nitrogen and sugar composition of ingested phloem sap and excreted honeydew. *Journal of Experimental Biology* 35, 74–84.
- Molyneux, R.J., Campbell, B.C., Dreyer, D.L., 1990. Honeydew analysis for detecting phloem transport of plant natural products. Implications for host-plant resistance to sap-sucking insects. *Journal of Chemical Ecology* 16, 1899–1909.
- Moser, S.E., Harwood, J.D., Obrycki, J.J., 2008. Larval feeding on Bt hybrid and non-Bt corn seedlings by *Harmonia axyridis* (Coleoptera: Coccinellidae) and *Coleomegilla maculata* (Coleoptera: Coccinellidae). *Environmental Entomology* 37, 525–533.
- Mueller, U.G., Schultz, T.R., Currie, C.R., Adams, R.M.M., Malloch, D., 2001. The origin of the attine ant-fungus mutualism. *Quarterly Review of Biology* 76, 169–197.
- Nalepa, C.A., Bambara, S.B., Burroughs, A.M., 1992. Pollen and nectar feeding by *Chilocorus kuwanae* (Silvestri) (Coleoptera: Coccinellidae). *Proceedings of the Entomological Society of Washington* 94, 596–597.
- Nedved, O., Ceryngier, P., Hodkova, M., Hodek, I., 2001. Flight potential and oxygen uptake during early dormancy in *Coccinella septempunctata*. *Entomologia Experimentalis et Applicata* 99, 371–380.
- Nijima, K., Abe, W., Matsuka, M., 1997. Development of low-cost and labor-saving artificial diet for mass production of an aphidophagous coccinellid, *Harmonia axyridis* (Pallas). *Bulletin of the Faculty of Agriculture, Tamagawa University* 37, 63–74.
- Omkar, 2006. Suitability of different foods for a generalist ladybird, *Micraspis discolor* (Coleoptera: Coccinellidae). *International Journal of Tropical Insect Science* 26, 35–40.
- Osenberg, C.W., Sarnelle, O., Goldberg, D.E., 1999. Meta-analysis in ecology: concepts, statistics, and applications. *Ecology* 80, 1103–1104.
- Pemberton, R.W., Vandenberg, N.J., 1993. Extrafloral nectar feeding by ladybird beetles (Coleoptera: Coccinellidae). *Proceedings of the Entomological Society of Washington* 95, 139–151.
- Percival, M.S., 1961. Types of nectar in angiosperms. *New Phytologist* 60, 235–281.
- Petanidou, T., 2005. Sugars in Mediterranean floral nectars: an ecological and evolutionary approach. *Journal of Chemical Ecology* 31, 1065–1088.
- Petanidou, T., Van Laere, A., Ellis, W.N., Smets, E., 2006. What shapes amino acid and sugar composition in Mediterranean floral nectars? *Oikos* 115, 155–169.
- Petanidou, T., Van Laere, A.J., Smets, E., 1996. Change in floral nectar components from fresh to senescent flower of *Capparis spinosa* (Capparidaceae), a nocturnally flowering Mediterranean shrub. *Plant Systematics and Evolution* 199, 79–92.
- Pilcher, C.D., Obrycki, J.J., Rice, M.E., Lewis, L.C., 1997. Preimaginal development, survival, and field abundance of insect predators on transgenic *Bacillus thuringiensis* corn. *Environmental Entomology* 26, 446–454.
- Putman, W.L., 1955. Bionomics of *Stethorus punctillum* Weise (Coleoptera: Coccinellidae) in Ontario. *Canadian Entomologist* 87, 9–33.
- Putman, W.L., 1963. Nectar of peach leaf glands as insect food. *Canadian Entomologist* 95, 108–109.
- Putman, W.L., 1964. Occurrence and food of some coccinellids (Coleoptera) in Ontario peach orchards. *Canadian Entomologist* 96, 1149–1155.
- Reznik, S.Y., Vaghina, N.P., 2006. Dynamics of fat content during induction and termination of “trophic diapause” in *Harmonia sedecimnotata* Fabr. females (Coleoptera, Coccinellidae). *Entomological Review* 86, 125–132.
- Ricci, C., Fiori, G., Colazza, S., 1983. Regime alimentare dell'adulto di *Tytthaspis sedecimpunctata* (L.) (Coleoptera: Coccinellidae) in ambiente a influenza antropica primaria: prato polifita. *Atti XIII Congr. Naz. Ital. Entomol., Sestriere-Torino*, 691–698.
- Ricci, C., Ponti, L., Pires, A., 2005. Migratory flights and pre-diapause feeding of *Coccinella septempunctata* (Coleoptera: Coccinellidae) adults in agricultural and mountain ecosystems of Central Italy. *European Journal of Entomology* 102, 531–538.
- Richards, D.R., Evans, E.W., 1998. Reproductive responses of aphidophagous lady beetles (Coleoptera: Coccinellidae) to nonaphid diets: an example from alfalfa. *Annals of the Entomological Society of America* 91, 632–640.
- Riddick, E.W., Barbosa, P., 1998. Impact of Cry3A-intoxicated *Leptinotarsa decemlineata* (Coleoptera: Chrysomelidae) and pollen on consumption, development, and fecundity of *Coleomegilla maculata* (Coleoptera: Coccinellidae). *Annals of the Entomological Society of America* 91, 303–307.
- Rockwood, L.P., 1952. Notes on coccinellids in the Pacific northwest. *Pan-Pacific Entomologist* 28, 139–147.
- Rosenberg, M.S., Adams, D.C., Gurevitch, J., 2000. *MetaWin Version 2*. Sinauer Associates, Sunderland, MA.
- Roulston, T.A., Buchmann, S.L., 2000. A phylogenetic reconsideration of the pollen starch-pollination correlation. *Evolutionary Ecology Research* 2, 627–643.
- Roulston, T.A., Cane, J.H., 2000. Pollen nutritional content and digestibility for animals. *Plant Systematics and Evolution* 222, 187–209.
- Roulston, T.A., Cane, J.H., Buchmann, S.L., 2000. What governs protein content of pollen: pollinator preferences, pollen-pistil interactions, or phylogeny? *Ecological Monographs* 70, 617–643.
- Shuel, R.W., 1955. Nectar secretion. *American Bee Journal* 95, 229–234.
- Seagraves, M.P., 2009. Lady beetle oviposition behavior in response to the trophic environment. *Biological Control* 51, 313–322.
- Simanton, F.L., 1916. *Hyperaspis binotata*, a predatory enemy of the terrapin scale. *Journal of Agricultural Research* 6, 197–203.
- Smith, B.C., 1961. Results of rearing some coccinellid (Coleoptera: Coccinellidae) larvae on various pollens. *Proceedings of the Entomological Society of Ontario* 91, 270–271.
- Smith, B.C., 1965. Growth and development of coccinellid larvae on dry foods (Coleoptera: Coccinellidae). *Canadian Entomologist* 97, 760–768.
- Smith, B.C., 1971. Effects of various factors on the local distribution and density of coccinellid adults on corn (Coleoptera: Coccinellidae). *Canadian Entomologist* 103, 1115–1120.
- Smith, L.L., Lanza, J., Smith, G.C., 1990. Amino acid concentrations in extrafloral nectar of *Impatiens sultani* increase after simulated herbivory. *Ecology* 71, 107–115.
- Smith, S.F., Krischik, V.A., 1999. Effects of systemic imidacloprid on *Coleomegilla maculata* (Coleoptera: Coccinellidae). *Environmental Entomology* 28, 1189–1195.

- Solbreck, C., 1974. Maturation of post-hibernation flight behaviour in the coccinellid *Coleomegilla maculata* (DeGeer). *Oecologia* 17, 265–275.
- Spellman, B., Brown, M.W., Matthews, C.R., 2006. Effect of floral and extrafloral resources on predation of *Aphis spiraeicola* by *Harmonia axyridis* on apple. *BioControl* 51, 715–724.
- Standifer, L.N., 1967. A comparison of the protein quality of pollens for growth-stimulation of the hypopharyngeal glands and longevity of honey bees, *Apis mellifera* L. (Hymenoptera: Apidae). *Insectes Sociaux*, Paris 14, 415–426.
- Standifer, L.N., Devys, M., Barbier, M., 1968. Pollen sterols – a mass spectrographic survey. *Phytochemistry* 7, 1361–1365.
- Stanley, R.G., Linskins, H.F., 1974. *Pollen: Biology, Biochemistry, Management*. Springer-Verlag, New York.
- Stephenson, A.G., 1982. The role of extrafloral nectaries of *Catalpa speciosa* in limiting herbivory and increasing fruit production. *Ecology* 63, 663–669.
- Sutherland, A.M., Parrella, M.P., 2009. Mycophagy in Coccinellidae: review and synthesis. *Biological Control* 51, 284–293.
- Tedders, W.L., Schaefer, P.W., 1994. Release and establishment of *Harmonia axyridis* (Coleoptera: Coccinellidae) in the southeastern United States. *Entomological News* 105, 228–243.
- Tooker, J.F., Hanks, L.M., 2000. Flowering plant hosts of adult hymenopteran parasitoids of Central Illinois. *Annals of the Entomological Society of America* 93, 580–588.
- Torres, C., Galetto, L., 2002. Are nectar sugar composition and corolla tube length related to the diversity of insects that visit Asteraceae flowers? *Plant Biology* 4, 360–366.
- Triltsch, H., 1997. Gut contents in field sampled adults of *Coccinella septempunctata* (Col.: Coccinellidae). *Entomophaga* 42, 125–131.
- Triltsch, H., 1999. Food remains in the guts of *Coccinella septempunctata* (Coleoptera: Coccinellidae) adults and larvae. *European Journal of Entomology* 96, 355–364.
- van den Meiracker, R.A.F., Hammond, W.N.O., van Alphen, J.J.M., 1990. The role of kairomones in prey finding by *Diomus* sp. and *Exochomus* sp., two coccinellid predators of the cassava mealybug, *Phenacoccus manihoti*. *Entomologia Experimentalis et Applicata* 56, 209–217.
- Vattala, H.D., Wratten, S.D., Phillips, C.B., Wäckers, F.L., 2006. The influence of flower morphology and nectar quality on the longevity of a parasitoid biological control agent. *Biological Control* 39, 179–185.
- Wäckers, F.L., 2000. Do oligosaccharides reduce the suitability of honeydew for predators and parasitoids? A further facet to the function of insect-synthesized honeydew sugars. *Oikos* 90, 197–201.
- Wäckers, F.L., 2001. A comparison of nectar- and honeydew sugars with respect to their utilization by the hymenopteran parasitoid *Cotesia glomerata*. *Journal of Insect Physiology* 47, 1077–1084.
- Wäckers, F.L., 2005. Suitability of (extra-)floral nectar, pollen, and honeydew as insect food sources. In: Wäckers, F.L., van Rijn, P.C.J., Bruin, J. (Eds.), *Plant-Provided Food for Carnivorous Insects*. Cambridge University Press, Cambridge, UK, pp. 17–74.
- Wade, M.R., Zalucki, M.P., Wratten, S.D., Robinson, K.A., 2008. Conservation biological control of arthropods using artificial food sprays: current status and future challenges. *Biological Control* 45, 185–199.
- Watson, J.R., Thompson, W.L., 1933. Food habits of *Leis conformis* Boisd. (Chinese ladybeetle). *Florida Entomologist* 17, 27–29.
- Way, M.J., 1963. Mutualism between ants and honeydew-producing Homoptera. *Annual Review of Entomology* 8, 307–344.
- Williams, L.L., Roane, T.M., Beach, J.P., 2005. Gustatory acceptance, longevity, and utilization of nectar and honeydew sugars by *Anaphes iole*, an egg parasitoid of *Lygus* bugs. In: Hoddle, M.S. (Ed.), *International Symposium on Biological Control of Arthropods*. United States Forest Service, Davos, Switzerland, pp. 409–419.
- Wolfenbarger, L.L., Naranjo, S.E., Lundgren, J.G., Bitzer, R.J., Watrud, L.S., 2008. Bt Crop effects on functional guilds of non-target arthropods: a meta-analysis. *PLoS One* 3, e2118. doi:10.1371/journal.pone.0002118.
- Woodring, J., Wiedemann, R., Volk, W., Hoffmann, K.H., 2006. Oligosaccharide synthesis regulates gut osmolality in the ant-attended aphid *Metopeurum fuscoviride* but not in the unattended aphid, *Macrosiphoniella tanacetaria*. *Journal of Applied Entomology* 131, 1–7.
- Yao, L., Akimoto, S.-I., 2002. Flexibility in the composition and concentration of amino acids in honeydew of the drepanosiphid aphid *Tuberculatus quercicola*. *Ecological Entomology* 27, 745–752.
- Yinon, U., 1969. Food consumption of the armored scale lady-beetle *Chilocoris bipustulatus* (Coccinellidae). *Entomologia Experimentalis et Applicata* 12, 139–146.